
PicoPhi RCA Premix Kit with Random Primers (2x)

Cat#: PM100 or PM1000



FOR RESEARCH USE ONLY

PicoPhi RCA Premix Kit is intended for molecular biology use and *in vitro* use only. This product is not intended for diagnosis, prevention or treatment of a disease in human beings or animals.

Store Kit at -80°C on Receipt



Description

PicoPhi RCA Premix Kit (Cat#: PM100) is a 2x premix version of PicoPhi RCA kit (Phi100), containing all components necessary for a successful and reliable **Rolling Circle Amplification (RCA)**, such as *PicoPhi* DNA Polymerase, dNTPs, and random primers etc, except circular DNA template. This premixed formulation saves time and reduces contamination due to a reduced number of pipetting steps required for RCA set up. The premix is optimized for efficient and reproducible RCA, an equivalent of *PicoPhi* DNA Amplification Kit (Cat#: Phi100).

Kit Components:

Components	100 Reactions (Cat#: PM100)	1000 Reaction (Cat#: PM1000)	Storage
Sample Buffer	800 µl	10x 800 µl	-20°C
PicoPhi Premix	500 µl	10x 500 µl	-20°C

Detailed Protocol

PicoPhi RCA Pre-mix Kit has capacity to detect as low as 1 pictogram of template DNA under optimal condition. 0.2~0.5 µl of saturated overnight culture or 1/10 to 1/100 of the colony (approximately $10^2 \sim 10^4$ cells) would be enough for PicoPhi RCA reaction. Please keep in mind that excess culture or colony will inhibit PicoPhi RCA reaction! The protocol described below is a general protocol for amplifying circular DNA from various sources. Yields and kinetics may vary if crude or un-quantified samples are amplified. We recommend to consider a starting point for adapting your specific reaction. Procedures of *PicoPhi* RCA reaction were outlined below.

1. Preparation of Sample Mix:

Sample Mix could be prepared, depending on material sources, as described below:

1.1. **Purified DNA or DNA ligation/assembly reactions:**

Transfer 4 µl of Sample Buffer into a 0.2 ml PCR tube.

Add 1 µl of circular DNA (≥ 1 pg/µl) to the above PCR tube.

Heat to 95 °C for 3 minutes and then quickly cool to 4°C.

Keep the samples on ice until use.

1.2. **Bacterial colonies:**

Transfer 5 µl of Sample Buffer into a 0.2 ml PCR tube.

Pick 1/10 to 1/100 of the colony (approximately $10^2 \sim 10^4$ cells) and add to the above PCR Tub.

Heat to 95 °C for 3 minutes and then quickly cool to 4°C.

Keep the samples on ice until use.

1.3. **Liquid bacterial culture:**

Transfer 4.0~4.8 µl of Sample Buffer into a 0.2 ml PCR tube.

Add 0.2~1.0 µl of saturated overnight culture to the above PCR tube.

Heat to 95 °C for 3 minutes and then quickly cool to 4°C.

Keep the samples on ice until use.



1.4. Glycerol stock:

Transfer 4.0~4.8 µl of Sample Buffer into a 0.2 ml PCR tube.

Add 0.2~1.0 µl of glycerol stock to the above PCR tube.

Heat to 95 °C for 3 minutes and then quickly cool to 4 °C.

Keep the samples on ice until use.

***Note:** Heating at higher temperature or longer time may increase the probability of nicking target DNA and releasing host genomic DNA into cell lysis to the reaction, where the host genomic DNA will compete with the desired template DNA during amplification.*

2. DNA Amplification:

Add 5 µl of *PicoPhi* pre-mix to the above PCR tube contains 5 µl of Sample Mix as showed below:

Component	Volume/reaction
Sample Mix with denatured DNA	5 µl
PicoPhi Premix	5 µl
Final Volume	10 µl

Mix well and incubate at 30 °C for 3~24 hrs.

***Note:** Actually, mixing any equal volume of sample mix and FemtoPhi Pre-mix will work fine.*

3. Inactivate PicoPhi: Incubating at 65 °C for 10 min, and then cool to 4 °C.

4. Perform Downstream Application:

- The amplified DNA can be directly used for the cycle sequencing reaction without purification;
- The amplified DNA can be directly used for DNA restriction enzyme digestion;
- An aliquot of the amplified DNA can be examined by agarose gel;
- Specially, the amplified RCA products as concatemers, maybe after diluted, can be used to directly transform *Bacillus*. Compared to the traditional *bacillus* transformation methods, the RCA1.0 products would give the highest transformation rate, which is of critical importance to *bacillus* gene cloning, expression and library construction.

5. FAQ and Troubleshooting: Please contact us at info@genexp-llc.com.

Limited Use License and Warranty

Use of this Product is subject to the following terms and conditions. Genxp LLC ("We") warrant that the Product meets the specifications described in this manual. If the terms and conditions are not acceptable, please return all components of Products with original receipt by the original buyer to Genxp LLC within 10 business days. This product is for internal research purposes only and is not for use in commercial applications of any kind, including, without limitation, quality control and commercial services such as reporting the results of purchaser's activities for a fee or other form of consideration. No right or license to resell this product or any of its components is conveyed expressly, by implication, or by estoppel. This Product should be used in accordance with the NIH guidelines developed for recombinant DNA and genetic research. Genxp LLC disclaims any and all responsibility for injury or damage which may be caused by the failure of the buyer or any other person to use the Product in accordance with the terms and conditions outlined herein. If you should have any questions or concerns about any of our products, please contact us at info@genexp-llc.com.